# RESEARCH FACTS





# IN PROGRESS

# Developing new sainfoin varieties

### **PROJECT TITLE**

Novel sainfoin cultivars for enhancing production efficiencies of pasture and beef cattle, and building capacity in forage breeding (BCRC project code: FRG.02.17)

### In Progress:

Results expected in March 2022

#### **RESEARCHERS**

Dr. Surya Acharya (PhD), Agriculture and Agri-Food Canada Lethbridge <a href="mailto:surya.acharya@agr.gc.ca">surya.acharya@agr.gc.ca</a>

Dr. Bill Biligetu (PhD), College of Agriculture and Bioresources, University of Saskatchewan bill.biligetu@usask.ca

Dr. Doug Cattani (PhD), University of Manitoba doug.cattani@umanitoba.ca

Agriculture and Agri-Food Canada Lethbridge: Dr. Tim McAllister (PhD), Dr. Yuxi Wang (PhD); AAFC Edmonton: Darren Bruhjell; AAFC Regina: Bill Houston

#### **Background:**

Grassstands are less productive than they could be because bloat fears discourage some producers from seeding alfalfa. Sainfoin is a non-bloating legume. It has lower protein levels than alfalfa, but contains condensed tannins that appear to reduce protein breakdown in the rumen and improve protein digestion and absorption in the intestine. Consequently, gains on sainfoin pasture can be as efficient and rapid as on alfalfa pasture. Sainfoin is resistant to the alfalfa weevil and grows earlier in the spring and later in the fall than alfalfa. Researchers at AAFC Lethbridge have been selecting sainfoin for improved yield, regrowth and survival in alfalfa stands and have found that sainfoin's survival depends partly on the alfalfa variety it is grown with as well as where it is grown.

#### **Objectives:**

To gain insight into grass and sainfoin interactions and to develop new sainfoin germplasm for grass/sainfoin mixed stands. Researchers will also determine nutritional quality of new sainfoin and grass/sainfoin stands and develop integrated crop management practice to optimize growth and longevity of grass/sainfoin pastures.

#### **What They Will Do:**

Five sainfoin populations will be seeded (within, between or across rows) with orchard, meadow and hybrid brome in Lethbridge, Saskatoon and Carman. Plants will be counted four weeks after seeding and yield and grass:sainfoin proportions will be measured twice per year for four years.

Forage quality will be monitored in years two to four with laboratory and animal digestibility trials using samples collected at times relevant to grazing and silaging.

The sainfoin plants that flourish the best in grass stands will be identified in years three and four, cloned and screened for disease resistance, seed production and forage quality, and other economically and agronomically relevant traits..

#### **Implications:**

This project will generate novel bloat-free sainfoin cultivars that improve the efficiencies of pasture and beef cattle production in Canada.

### Proudly funded by:







Factsheet information provided by the Beef Cattle Research Council

beefresearch.ca